Nutrient Broth

Intended Use

Nutrient Broth is used for the cultivation of many species of nonfastidious microorganisms.

Summary and Explanation

Nutrient Broth has the formula originally designed for use in the *Standard Methods for Examination of Water and Waste-water*. It is not a recommended bacteriological medium in later editions of this publication. It is one of several nonselective media useful in routine cultivation of microorganisms.¹⁻³

Principles of the Procedure

This relatively simple formulation supports the growth of non-fastidious microorganisms due to its content of peptone and beef extract.

User Quality Control

Identity Specifications **Difco™ Nutrient Broth**

Dehydrated Appearance: Medium tan, free-flowing, homogeneous.

Solution: 0.8% solution, soluble in purified water. Solu-

tion is light to medium amber, clear.

Prepared Appearance: Light to medium amber, clear

Reaction of 0.8%

Solution at 25°C: pH 6.8 ± 0.2

Cultural Response Difco™ Nutrient Broth

Prepare the medium per label directions. Inoculate and incubate at $35 \pm 2^{\circ}$ C for 18-24 hours.

ORGANISM	ATCC™	INOCULUM CFU	RECOVERY
Escherichia coli	25922	$10^2 - 10^3$	Good
Staphylococcus aureus	25923	10 ² -10 ³	Good

Formula

Difco™ Nutrient Broth

Approximate Formula* Per Liter	
Beef Extract3.0	g
Peptone	g
*Δdiusted and/or supplemented as required to meet performance criteria	

Directions for Preparation from Dehydrated Product

- 1. Dissolve 8 g of the powder in 1 L of purified water.
- 2. Autoclave at 121°C for 15 minutes.
- 3. Test samples of the finished product for performance using stable, typical control cultures.

Procedure

Inoculate tubes of the broth medium with the test samples. Incubate tubes for 18-24 hours at 35 ± 2 °C in an aerobic atmosphere.

Expected Results

After incubation, growth is evidenced by the appearance of turbidity in the broth. Aliquots of the broth can be used for subculturing to solid media for purification and identification purposes.

References

- Marshall (ed.). 1993. Standard methods for the examination of dairy products, 16th ed. American Public Health Association, Washington, D.C.
- Public Health Association, Washington, D.C.
 2. U.S. Food and Drug Administration. 2001. Bacteriological analytical manual, online. AOAC International, Gaithersburg, Md.
- tional, Gaithersburg, Md.

 3. Downes and Ito (ed.). 2001. Compendium of methods for the microbiological examination of foods, 4th ed. American Public Health Association, Washington, D.C.

Availability

Difco™ Nutrient Broth

AOAC	BAM CCAI	M COMPF SMD
Cat. No.	233000	Dehydrated – 100 g
	234000	Dehydrated – 500 g
	231000	Dehydrated – 2 kg
	232000	Dehydrated – 10 kg

BBL™ Nutrient Broth

BAM CCAM COMPF SMD

Cat. No. 221669 Prepared Tubes, 5 mL (K Tubes) – Pkg. of 10

