
Catheter Care Awareness Week puzzles and games

Preventing Catheter-Associated Urinary Tract Infections crossword puzzle



Across

- The CDC recommends ______ precautions when touching or manipulating an indwelling catheter, including gloves and a gown, if necessary
- Urinary retention management may include the use of intermittent catheters and bladder
- This technique is recommended for inserting an indwelling catheter and includes the use of sterile supplies, such as gloves, drapes, single-use lubricant packets, and sterile or antiseptic solution for periurethral cleaning
- When a patient no longer meets indications for an indwelling catheter, prompt ______ is recommended to reduce the risk of infection
- 5. _____ catheterization and bladder scanning may be recommended for patients with postoperative urinary retention.

Down

- This type of catheter has a curved tip and is used to help navigate obstructions in the urethra, such as enlarged prostate
- Avoiding kinks and ______ loops in catheter tubing helps reduce the risk of urinary stasis or backflow of urine, which can increase the risk of infection
- The CDC recommends an indwelling catheter has proper ____ to prevent movement and urethral traction
- 4. Supplies used for inserting an indwelling catheter should be
- 5. Urinary catheter _____ is the biggest risk factor for a catheterassociated urinary tract infection (CAUTI)

Α	Т	U	В	L	W	R	1	Р	D	Α	W	С	- 1	N	G	Е
Р	В	Υ	Α	N	G	W	Α	R	U	С	I	R	Е	Т	L	Х
В	L	L	Υ	Н	В	I	М	Е	Q	U	С	W	F	S	В	Е
Р	С	٧	Α	Р	D	0	Р	Α	D	S	K	Е	S		Υ	D
S	Т	Н	K	D	Р	U	С	Α	Р	D	I	J	L	S	Н	L
В	Х	С	М	R	D	G	K	U	М	Н	N	С	Α	В	Z	Р
С	Е	Т	0	ı	L	Е	Т	1	N	G	G	0	N	N	U	S
0	0	D	Α	D	D	ı	R	ı	F	Α	D	U	I	Т	R	0
М	S	В	Р	Z	Е	W	J	S	N	С	N	F	R	Т	L	Х
М	W	С	S	Α	В	L	ı	D	С	G	L	Е	U	Р	N	Н
0	Е	0	М	Κ	N	S	Α	С	0	Α	Н	R	0	Е	Α	L
D	U	М	S	N	F	U	R	G	ı	W	N	U	Т	ı	В	М
Е	0	Т	N	Е	Т	Т	1	М	R	Е	Т	N	I	N	N	М
R	K	Е	Х	٧	J	W	Е	Н	М	٧	I	Υ	Е	Т	Z	G
R	٧	Т	Е	Χ	Т	Е	R	N	Α	L	S	Χ	Z	R	R	Т
Т	В	М	G	0	С	Т	L	D	D	Α	Н	L	С	U	1	Υ
K	٧	0	Х	F	R	0	R	K	Α	N	J	R	Α	Т	S	0

Alternatives to indwelling catheters word search

EXTERNALS
INTERMITTENT
BLADDER SCANNER
BEDPAN
URINALS
COMMODE
TOILETING
WICKING
PADS







What would you change in the maintenance of these catheters?









Alternatives to indwelling catheters word search answers

Α	Т	U	В	L	W	R	1	Р	D	Α	W	С	1	N	G	Е
Р	В	Υ	Α	N	G	W	Α	R	U	С	1	R	Е	Т	L	Х
В	L	L	Υ	Н	В	1	М	Е	Q	U	С	W	F	S	В	Е
Р	С	٧	Α	Р	D	0	Р	Α	D	S	K	Ε	S	- 1	Υ	D
S	Т	Н	K	D	Р	U	С	Α	Р	D	-1	J	L	S	Н	L
В	Х	С	М	R	D	G	K	U	М	Н	N	С	Α	В	Z	Р
С	Е	Т	0	-1	L	Е	Т	1	N	G	G	0	N	N	U	S
О	0	D	Α	D	D	-1	R	- 1	F	Α	D	U	-1	Т	R	0
М	S	В	Р	Z	Е	W	J	S	N	С	N	F	R	Т	L	Х
М	W	С	S	Α	В	L	1	D	С	G	L	Е	U	Р	N	Н
0	Е	0	М	K	N	S	Α	С	0	Α	Н	R	0	Е	Α	L
D	U	М	S	N	F	U	R	G	-1	W	N	U	Т	-1	В	М
Е	0	Т	N	Е	T	Т	1	М	R	Е	Т	N	-1	N	N	М
R	K	Е	Х	٧	J	W	Е	Н	М	٧	I	Υ	Е	Т	Z	G
R	٧	Т	Е	Х	T	Е	R	N	Α	L	S	Х	Z	R	R	Т
T	В	М	G	0	С	Т	L	D	D	Α	Н	L	С	U	-1	Υ
K	٧	0	Х	F	R	0	R	K	Α	N	J	R	Α	Т	S	0

Crossword puzzle answer key

D.

Across

- The CDC recommends STANDARD precautions when touching or manipulating an indwelling catheter, including gloves
- Urinary retention management may include the use of intermittent catheters and bladder SCANNERS
- This technique is recommended for inserting an indwelling catheter and includes the use of sterile supplies, such as gloves, drapes, single-use lubricant packets, and sterile or antiseptic solution for periurethral cleaning ASEPTIC
- When a patient no longer meets indications for an indwelling catheter, prompt REMOVAL is recommended to reduce the risk of infection
- INTERMITTENT catheterization and bladder scanning may be recommended for patients with postoperative urinary

Down

- This type of catheter has a curved tip and is used to help navigate obstructions in the urethra, such as enlarged prostate **COUDÉ**
- Avoiding kinks and **DEPENDENT** loops in catheter tubing helps reduce the risk of urinary stasis or backflow of urine, which can increase the risk of infection
- The CDC recommends an indwelling catheter has proper ${f SECUREMENT}$ to prevent movement and urethral traction
- Supplies used for inserting an indwelling catheter should be STERILE
- Urinary catheter **DURATION** is the biggest risk factor for a catheter-associated urinary tract infection (CAUTI)

What Would You Change in the Maintenance of These Catheters?

- D. Dependent loop; no securement device; broken system/no tamper evident seal
- B. Bag on floor; no securement device; broken system/no tamper evident seal in place and above bladder; dependent loop; no securement device; broken system/no tamer evident seal in place.

Dependent loop; no securement device

BD[®] Clinical Connect Product Training and Education Program



bd.com