



Victoria Nurse

Using the needle-free blood collection device to enhance the experience of patients.^{1*}

The hospital floor Victoria works on has a mix of some very sick patients and some more stable patients who are able to leave the hospital after only a few days. No matter the condition, one procedure that remains tough across patients is blood draws.

24% & **63%**
of adults of children
are needle phobic.²

Victoria was able to use the PIVO™ Needle-free Blood Collection Device for the first time with her patient, Wyatt.

“Using the device was super easy.”

Victoria was surprised how well the procedure went, both for her and for Wyatt. She remarked, “I guess I was expecting to have to troubleshoot but everything went smoothly.”

“The patient didn’t experience discomfort. He didn’t even know what I was doing.”

Learn how the PIVO™ Needle-free Blood Collection Device may advance your practice at bd.com/PIVO



* Compared to traditional blood draws



The BD Peripheral Line Draw Solution

Trust in the PIVO™ Needle-free Blood Collection Device and the ExT™ Stabilized Extension Set with NearPort™ IV Access, the practice-advancing, needle-free blood draw solution. Elevate the standard of care while instilling confidence and enhancing experience for both your patients and clinicians.¹



Reliable

Reduce the risk of preanalytical lab errors and sample re-collects.³ Improve confidence in diagnostic accuracy, helping your patients receive the right treatment at the right time.³



Safe

Helps protect patients by reducing risks of IV replacements and is an alternative to central line draws while aiding in vessel health preservation.^{3,4,5}



Comfort

Help alleviate the fear associated with repetitive needlesticks.⁶ Draw blood while minimizing disruption to your patients' rest and healing processes.^{6,7}

When compared to venipuncture, patients who received daily phlebotomy preferred the PIVO™ Needle-free Blood Collection Device experience.⁶

9.1 out of 10

Learn how the PIVO™ Needle-free Blood Collection Device may advance your practice at bd.com/PIVO

BD, Sandy, UT, 84070, U.S.

References

- 1 Cadacio C, Nachamkin I. A novel needle-free blood draw device for sample collection from short peripheral catheters. *JJN*. 2017;40(3):156-162.
- 2 Taddio A, Ipp M, Thivakaran S, et al. Survey of the prevalence of immunization non-compliance due to needle fears in children and adults. *Vaccine*. 2012;30(32):4807-4812. doi:10.1016/j.vaccine.2012.05.011.
- 3 Pendleton B, LaFaye R. Multicenter study of needle-free blood collection system for reducing specimen error and intravenous catheter replacement. *J Healthc Qual*. 2022;44(2):e24-e30. doi:10.1097/JHQ.0000000000000331
- 4 Kuriakose L. Decreasing central line associated bloodstream infection through limiting the use of central venous catheters for routine blood draws. *J Dr Nurs Pract*. 2020;13(2):173-183. doi:10.1891/JDNP-D-19-00071
- 5 O'Malley C. The frequency and reasons for central line accesses in critical care units. *Canadian Journal of Infection Control*. 2018;33(3):165-167.
- 6 Mulloy DF, Lee SM, Gregas M, Hoffman KE, Ashley SW. Effect of peripheral IV based blood collection on catheter dwell time, blood collection, and patient response. *Appl Nurs Res*. 2018;40:76-79. doi:10.1016/j.apnr.2017.12.006
- 7 Matthews EE. Sleep disturbances and fatigue in critically ill patients. *AACN Adv Crit Care*. 2011;22(3):204-224. doi:10.1097/NC1.0b013e31822052cb

