


ChloraPrep skin antiseptis system

2% Chlorhexidine Gluconate w/v and 70% Isopropyl Alcohol v/v
Licenced, sterile, ANTT[®], latex free, controlled flow



Pinch

Hold the applicator as shown, being careful not to touch the sponge.

Pinch the lever to release the solution. You will hear a 'pop' as the ampoule breaks.

3ml with tint



NHS supply chain
order code:
MRB494



Apply

Starting at the incision site, gently press the applicator against the skin until the solution soaks the sponge.

Apply using repeated **up and down, back and forth strokes for at least 30 seconds**, before working outwards to the periphery.

10.5ml with tint



NHS supply chain
order code:
MRB495



Dry

Leave the area to air dry completely before applying sterile drapes. Do not blot or wipe away. Discard the applicator after a single use.

Important safety point:
Do not drape or use ignition source until the solution has completely dried.

26ml with tint



NHS supply chain
order code:
MRB496

For customer services and all other enquiries:

Please telephone: 0800 917 8776

Email: uk-customer-service@carefusion.com

or visit: chloraprep.co.uk

Prescribing information on reverse



Prescribing Information:

ChloroPrep™ & ChloroPrep with Tint 2% chlorhexidine gluconate w/v / 70% isopropyl alcohol v/v cutaneous solution. Refer to the Summary of Product Characteristics before prescribing. Presentation: ChloroPrep – each applicator contains 0.67 mL, 1.5 mL, 3 mL, 10.5 mL or 26 mL of 20 mg/mL chlorhexidine & 0.70 mL/mL isopropyl alcohol; ChloroPrep with Tint – each applicator contains 3 mL, 10.5 mL or 26 mL of 20 mg/mL chlorhexidine & 0.70 mL/mL isopropyl alcohol. Indication: Disinfection of skin prior to invasive medical procedures. Dosage & administration: Applicator volume dependent on invasive procedure being undertaken. May be used for all age groups and patient populations. Use with care in newborn babies and those born prematurely. Applicator squeezed to break ampoule and release antiseptic solution onto sponge. Solution applied by gently pressing sponge against skin and moving back and forth for 30 seconds. The area covered should be allowed to air dry. Contraindications: Patients with known hypersensitivity to ChloroPrep or ChloroPrep with Tint or any of its components, especially those with a history of possible Chlorhexidine-related allergic reactions. Warnings and precautions: Solution is flammable. Do not use with ignition sources until dry. Do not use in excessive quantities, allow to pool in patient skin folds or drip on materials in contact with patient skin. Remove any soaked materials before proceeding with the intervention. Ensure no excess product is present prior to application of occlusive dressing. For external use only on intact skin, do not use on open skin wounds or broken or damaged skin. Over-vigorous use on fragile or sensitive skin or repeated use may lead to local skin reactions. Avoid prolonged skin contact. Avoid contact with eyes, mucous membranes, middle ear and neural

tissue. Chlorhexidine may induce hypersensitivity, including generalised allergic reactions and anaphylactic shock. May cause chemical burns in neonates, with a higher risk in preterm infants and within the first 2 weeks of life. Pregnancy & lactation: Although no studies have been conducted, no effects are anticipated as systemic exposure is negligible. Undesirable effects: Very rare; allergic or irritation skin reactions to chlorhexidine, isopropyl alcohol or sunset yellow (E110, present in ChloroPrep with Tint only), including erythema, rash, pruritus and blisters or application site vesicles, other local symptoms have included skin burning sensation, pain and inflammation. Frequency not known; hypersensitivity including anaphylactic shock, dermatitis, eczema, urticaria, chemical burns in neonates. Discontinue use at the first sign of local skin reaction. Per applicator costs (ex VAT) ChloroPrep: 0.67 mL (Sepp) - 30p; 1.5 mL (Frepp) - 55p; 1.5 mL – 78p; 3 mL – 85p; 10.5 mL - £2.92; 26 mL - £6.50. ChloroPrep with Tint: 3 mL – 89p; 10.5 mL – £3.07; 26 mL - £6.83 Legal category: GSL Marketing Authorisation Numbers: ChloroPrep, PL31760/0004 & ChloroPrep with Tint, PL31760-0001 Marketing Authorisation Holder: CareFusion UK 244 Ltd, The Crescent, Jays Close, Basingstoke, Hampshire, RG22 4BS. Date of Preparation: February 2016

Reporting suspected adverse reactions is important to monitor the benefit/risk balance of the medicinal product. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard. Adverse events should also be reported to CareFusion Freephone number: 0800 0437 546 or email: CareFusionGB@professionalinformation.co.uk

BD UK, Basingstoke, HPH, RG22 4BS, UK
0800 917 8776

bd.com

CareFusion is a wholly owned subsidiary of Becton, Dickinson and Company.
© 2016 BD. BD, the BD Logo and ChloroPrep are trademarks of Becton, Dickinson and Company. 0000CF02926 Issue 1

