

BLOOD CULTURE PREP KIT II

Cat. No. 260307

NDC 54365-200-99



1 FREPP®

**70% isopropyl alcohol
Antiseptic**

0.04 fl. oz. (1.1 ml)



1 SEPP®

**iodine tincture 2% USP
Patient Preoperative Skin
Preparation**

0.02 fl. oz. (0.67 ml)

CareFusion

El Paso, TX 79912

Made in USA

**NON-STERILE SOLUTION
APPLICATOR IS STERILE IF
PACKAGE IS INTACT**

STORE BELOW 30°C OR 86°F

FOR SINGLE &
EXTERNAL USE ONLY

FLAMMABLE, KEEP AWAY
FROM FLAME OR FIRE

DO NOT USE WITH
ELECTROCAUTERY
PROCEDURES



Bend and
Peel

BLOOD CULTURE PREP KIT II

Isopropyl alcohol 70% FREPP

Drug Facts

Active ingredient

Isopropyl alcohol 70% v/v, USP Antiseptic

Purpose

Antiseptic

Use for preparation of the skin prior to an injection

Warnings

For external use only. Flammable: Keep away from fire or flame

Do not use with electrocautery procedures

Do not use ■ in the eyes ■ on patients with known allergies to isopropyl alcohol

When using this product keep out of eyes, ears, and mouth

Stop use and ask a doctor if irritation, sensitization, allergic reaction, or other side effects occur. You may report side effects to FDA at 1-800-FDA-1088.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions ■ locate the vein to be used

■ remove Frepp from kit. Hold in a horizontal position and pinch handle ONCE to break ampule. Do not continue to squeeze handle.

■ place sponge on selected venipuncture site and depress once or twice to saturate sponge

■ scrub in a gentle back and forth motion for at least 30 seconds

■ allow to dry

Other information store below 30 °C (86 °F)

Inactive ingredient USP purified water

Questions? Call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

Iodine tincture 2% SEPP

Drug Facts

Active ingredient

Iodine 2% w/v, USP Antiseptic

Purpose

Antiseptic

Use patient preoperative skin preparation. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only.

Do not use on patients with known allergies to ethyl alcohol or iodine

When using this product keep out of eyes, ears, and mouth

Stop use and ask a doctor if irritation, sensitization, allergic reaction, or other side effects occur. You may report side effects to FDA at 1-800-FDA-1088.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions ■ remove the applicator wrapper. Do not touch the applicator tip.

■ with the applicator tip facing downward, pinch the barrel of the applicator ONCE to release the antiseptic.

■ press the applicator tip against the patient's skin. With gentle back and forth actions apply solution to the treatment area.

■ the area covered should be allowed to dry naturally.

■ proceed with collection of blood

■ discard after single use

Other information store below 30 °C (86 °F)

Inactive ingredients Ethyl alcohol, USP purified water, sodium iodide USP

Questions? Call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)